

IN THE CLAIMS

Please amend the claims as follows:

1. (Amended) A sweetener composition, comprising a mixture of:

(a) N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine-1-methyl ester; and

(b) Acesulfame K,

[wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by weight based on the total amount of said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame K] wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester comprises a C-type crystal or an A-type crystal.

12. (Amended) A drink composition, comprising:

(A) a mixture, comprising

(a) N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester;

and

(b) Acesulfame K;

and

(B) a potable liquid,

[wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by weight based on the total amount of said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame K] wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester comprises a C-type crystal or an A-type crystal.

13. (Amended) A method for preparing a sweetener composition, comprising:

(1) drying A-type crystals of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester, to obtain C-type crystals of

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester; and

(2) mixing said C-type crystals of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with Acesulfame K, to obtain a mixture,

[wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by weight based on the total weight of said

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame K] wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester comprises a C-type crystal having a water content of less than 3% by weight.

14. (Amended) A method for producing a sweetener, comprising:

(1) mixing N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with Acesulfame K, to obtain a mixture, [wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by weight based on the total weight of said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame K] wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester comprises a C-type crystal or an A-type crystal.

16. (Amended) A method for improving the dissolution rate of N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester, comprising:

(1) mixing said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester with Acesulfame K, prior to dissolving said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester, to obtain a mixture,

[wherein said Acesulfame K is present in said mixture in an amount of 50 to 97% by weight based on the total weight of said

N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester and said Acesulfame K] wherein said N-[N-(3,3-dimethylbutyl)-L- α -aspartyl]-L-phenylalanine 1-methyl ester comprises a C-type crystal or an A-type crystal.